Timor-Leste

Human Papillomavirus and Related Cancers, Fact Sheet 2017 (2017-04-19)



I. Key data on HPV and HPV-related cancers



Timor-Leste has a population of 337,309 women ages 15 years and older who are at risk of developing cervical cancer. Current estimates indicate that every year 46 women are diagnosed with cervical cancer and 24 die from the dis-

ease. Cervical cancer ranks as the 3rd most frequent cancer among women in Timor-Leste and the 2nd most frequent cancer among women between 15 and 44 years of age. Data is not yet available on the HPV burden in the general population of Timor-Leste. However, in South-Eastern Asia, the region Timor-Leste belongs to, about 3.0% of women in the general population are estimated to harbour cervical HPV-16/18 infection at a given time, and 71.6% of invasive cervical cancers are attributed to HPVs 16 or 18.

Table 1. Crude incidence rates of HPV-related cancers

	Male	Female
Cervical cancer	-	7.9
Anal cancer	-	-
Vulva cancer	-	-
Vaginal cancer	-	-
Penile cancer	-	-
Pharynx (excluding nasopharynx)	4.8	1.2


Table 2. Burden of cervical cancer

	Incidence	Mortality
Annual number of new cases/deaths	46	24
Crude rate	7.9	4.1
Age-standarized rate	13.3	8.1
Cumulative risk 0-74 years (%)	1.3	0.8
Ranking of cervical cancer (all years)	3rd	3rd
Ranking of cervical cancer (15-44 years)	2nd	8th

Table 3. Burden of cervical HPV infection Timor-

LUGIC		
	No. Tested	% (95% CI)
HPV prevalence in women with normal		
cytology	-	
HPV 16/18 prevalence:		
Normal cytology	-	
Low-grade cervical lesions	-	
High-grade cervical lesions	-	
Cervical cancer	-	

Figure 1. Comparison of the ten most frequent HPV oncogenic types in Timor-Leste among women with and without cervical lesions



*No data available. No more types than shown were tested or were positive.

Timor-Leste

HPV Information centre

Human Papillomavirus and Related Cancers, Fact Sheet 2017 (2017-04-19)

II. Complementary data on cervical cancer prevention

Table 4. Factors contributing to cervical cancer (cofactors)

Smoking prevalence (%), women	-
Total fertility rate (live births per women)	5.7
Hormonal contraception use (%) (pill, injectable or implant), among women	18.2
HIV prevalence (%), adults (15-49 years)	-

Table 7. Cervical screening practices and recommendations

Cervical cancer screening coverage, % (age and screening interval, reference)	-
Screening ages (years)	-
Screening interval (years) or frequency of screens	-

Table 5. Sexual behaviour

MEN	
Percentage of 15-year-old who have had sexual intercourse	-
Range of median age at first sexual intercourse	22.2-23.9
WOMEN	
Percentage of 15-year-old who have had sexual intercourse	-
Range of median age at first sexual intercourse	20.5-23.4

Figure 2. Estimated coverage of cervical cancer screening in Timor-Leste, by age and study




Table 6. HPV vaccine introduction

HPV vaccination programme	No Program
Date of HPV vaccination routine immunization programme start	-
HPV vaccination target age for routine immunization	-
Full course HPV vaccination coverage for routine immunization:	
% (calendar year)	-

Contact information:

ICO HPV Information Centre

Institut Català d'Oncologia

Avda. Gran Via de l'Hospitalet, 199-203

08908 L'Hospitalet de Llobregat (Barcelona, Spain)

e-mail: info@hpvcentre.net

internet adress: www.hpvcentre.net